

Knowledge, Attitudes and Practices of Healthcare Workers regarding Obesity: A Survey at a Tertiary Hospital in Vietnam

Tuan Doan Minh Tran

Department of Endocrinology and Nephrology, Le Van Thinh Hospital, Ho Chi Minh City, Vietnam

Abstract

Background. The global prevalence of obesity has tripled over the past four decades, with an estimated one-third of the world's population currently classified as overweight or obese. Obesity is recognized as one of the most urgent global public health challenges, with increasing rates observed in both developed and developing countries. In Southeast Asia, the burden of obesity is particularly concerning due to the coexistence of undernutrition and overnutrition, creating a double burden of disease. Healthcare workers play a critical role in the prevention, early detection and management of obesity. However, many healthcare workers lack updated knowledge on obesity, which is often associated with insufficient or inappropriate management.

Objective. To assess the knowledge, attitudes, and practices (KAP) of healthcare workers toward obesity management at a tertiary hospital in Ho Chi Minh City, Vietnam, and to identify associated barriers and interprofessional differences.

Methodology. A cross-sectional study was conducted among 232 healthcare workers using a 22-item structured, self-administered questionnaire. The survey evaluated knowledge of obesity risk factors and treatments, attitudes toward patient responsibility and stigma, and clinical practices such as screening, counseling and referrals. Associations between professional roles and responses were analyzed using Chi-square and Fisher's exact tests.

Results. Among 232 participants (37.5% were physicians and 62.5% were non-physician healthcare workers), only 63.8% [95% CI: 57.3–70.3%] recognized obesity as a chronic disease and fewer than half were aware of approved pharmacotherapies. While most workers believed they could assist patients, a significant majority (79.7% [95% CI: 75.0–84.9%]) attributed weight loss primarily to patient responsibility, indicating widespread implicit weight bias. Furthermore, physicians reported significantly higher odds of inadequate knowledge of weight management compared to other staff (OR = 4.5, 95% CI: 2.1–9.5).

Conclusion. Significant knowledge gaps and implicit weight biases persist among healthcare workers regarding obesity management. Addressing these educational and systemic barriers through specialized training and multidisciplinary care models in Vietnam can provide a crucial framework for improving obesity management across rapidly urbanizing Southeast Asian nations facing similar metabolic crises.

Key words: *obesity, healthcare worker, knowledge, attitudes, practice, Vietnam*

INTRODUCTION

Obesity has emerged as one of the most pressing global public health crises of the 21st century. The global prevalence of obesity has tripled over the past four decades, with an estimated one-third of the world's population currently classified as overweight or obese.¹ According to the updated World Health Organization (WHO) 2024 report, the global obesity rate has doubled between 1990 and 2022. By 2022, approximately 46% of the global adult

population was classified as overweight and 16% as obese.² In Asia, including Southeast Asia, the prevalence of obesity is projected to increase by 10%–20%.³

Despite this escalating crisis, a significant gap in clinical practice remains. Healthcare workers (HCWs) play a critical role in the prevention, early detection and management of obesity; however, many lack updated knowledge and formal training, potentially resulting in insufficient or inappropriate care.⁴

eISSN 2308-118x (Online)

Printed in the Philippines

Copyright © 2026 by Tran.

Received: January 23, 2026. Accepted: February 28, 2026.

Published online first: August 12, 2026.

<https://doi.org/10.15605/jafes.041.02.5583>

Corresponding author: Tuan Doan Minh Tran, MD

Deputy Head of Department

Le Van Thinh Hospital – Endocrinology and Nephrology Department,

130 Le Van Thinh St., Binh Trung Ward, Ho Chi Minh City, Vietnam, 700000

Tel. No. : (+84) 1900633878 loc. 6336

E-mail: trandoanminhtuan@gmail.com

ORCID: <https://orcid.org/0000-0002-5906-8611>

In Southeast Asia, the burden is particularly concerning due to the coexistence of undernutrition and overnutrition, creating a double burden of disease.⁵ Vietnam's rapid urbanization, marked by changes in dietary patterns and lifestyle behaviors and reductions in physical activity, has led to a sharp rise in overweight and obesity rates.^{6,7} Notably, higher obesity prevalence rates and metabolic risks are observed in this population when applying the Asian-specific BMI cutoffs (overweight ≥ 23 kg/m², obesity ≥ 25 kg/m²) compared to the standard WHO classification.⁸⁻¹¹ In 2022, the Vietnamese Ministry of Health issued national guidelines on the diagnosis and treatment of obesity,¹² reflecting growing attention from health authorities. However, obesity is still underrecognized in both society and the healthcare sector. Many healthcare workers lack updated knowledge on obesity, which may be linked to insufficient or inappropriate management.

Although previous studies have examined patient perceptions of obesity,¹³ there is little research that has focused on Healthcare workers. Therefore, assessing the knowledge, attitudes, and practices (KAP) of healthcare workers is essential. Such evaluation could inform strategies to improve professional training, reshape attitudes and enhance obesity management, thereby contributing to population-level efforts to reduce overweight and obesity rates. Based on these objectives, we conducted this study to investigate the healthcare workers' knowledge, attitudes and practices regarding obesity.

METHODOLOGY

Study design

This was a descriptive cross-sectional, multi-departmental study conducted at Le Van Thinh Hospital.

Study population

Sample size estimation

The minimum required sample size was estimated using the formula for a single population proportion:

$$n = Z_{[1-(\alpha/2)]}^2 \frac{p(1-p)}{d^2}$$

We set the confidence level at 95% ($Z = 1.96$) and the margin of error at 5% ($d = 0.05$). The expected proportion (p) was set to 0.84, based on a previous regional study across South and Southeast Asia in which 84% of healthcare workers correctly recognized obesity as a chronic disease.¹⁴ This calculation yielded a minimum sample size of 207. Our final recruited sample of 232 healthcare workers exceeded this requirement, ensuring adequate statistical power to address the research objectives and account for any potential nonresponse or incomplete data.

Participants were recruited using convenience sampling between June and September 2024. The research team introduced the study objectives and invited clinical staff to participate during regular departmental meetings. Inclusion criteria comprised clinical healthcare workers (physicians, nurses, midwives and technicians) actively working at the hospital for at least three months and provided written informed consent. Exclusion criteria included non-clinical administrative staff, individuals who declined to participate and those who submitted incomplete questionnaires. A total of 232 fully completed questionnaires were collected and retained for analysis. A paper-based questionnaire was administered to clinical staff during regular departmental meetings. The completion process, supervised by the research team to ensure data integrity, took approximately 15 minutes per participant.

Questionnaire development and validation

The 22-item questionnaire was developed based on the 2022 National Guidelines for the Diagnosis and Management of Obesity released by the Vietnamese Ministry of Health and adapted from previously validated KAP instruments reported in the recent literature. Face and content validity were evaluated by an expert panel comprising two endocrinologists and one public health specialist. Following expert revisions, the instrument was pilot-tested on a purposive sample of 20 healthcare workers (including physicians and nurses) to assess clarity, cultural appropriateness and completion time. To avoid prior exposure bias, these pilot participants were excluded from the final study sample. The internal consistency of the finalized questionnaire was evaluated using the full study sample ($N = 232$), demonstrating excellent reliability with an overall Cronbach's alpha of 0.850. The domain-specific alpha coefficients were 0.819 for Knowledge, 0.689 for Attitudes, and 0.646 for Practices.

Definition of outcomes and exposures

The primary outcomes were the proportions of healthcare workers demonstrating adequate knowledge, positive attitudes and appropriate clinical practices regarding obesity management. Secondary outcomes included the frequency of reported barriers to obesity management and the interprofessional differences in KAP domains.

Composite scores were not generated; KAP domains were analyzed individually. Knowledge items were categorized based on the Vietnamese Ministry of Health Guidelines. Attitudes were evaluated as positive if participants agreed with evidence-based statements regarding obesity care. Practices were assessed based on the appropriateness of clinical recommendations (e.g., lifestyle modifications, pharmacotherapy or referral) across different patient body mass index (BMI) and comorbidity profiles.

The primary exposures evaluated for their association with KAP outcomes were participants' professional roles

(categorized as physicians versus other healthcare workers, including nurses, midwives and technicians), BMI and years of medical experience.

Data analysis

Data completeness was verified prior to entry. Questionnaires with missing responses were excluded. Only fully completed forms were retained for the final analysis, ensuring no missing data within the analyzed dataset.

Data were entered using Epidata version 3.1 and analyzed with SPSS version 26.0. Before inferential analysis, the normality of continuous variables was assessed using the Shapiro-Wilk test. Normally distributed data were summarized using means and standard deviations, whereas non-normally distributed data were presented as medians and interquartile ranges (IQR). Descriptive statistics were used to present the data as percentages.

Associations between categorical variables were evaluated using the Pearson Chi-square test. The assumptions for Chi-square testing were verified prior to interpretation; Fisher's exact test was used where more than 20% of expected cell frequencies were less than 5. The threshold for statistical significance was defined as a two-tailed p -value of <0.05 . To adjust for potential confounders, stratified analyses by participant BMI and years of medical experience were performed to assess their association with the KAP outcomes.

Ethical considerations

Participation in this study was strictly voluntary. Participants were explicitly informed that they could

decline to participate or withdraw from the study at any time without negative consequences to their employment, performance evaluations or professional standing at the hospital.

To ensure absolute participant confidentiality, the study was designed to be completely anonymous. We did not collect any personally identifiable information, such as names, employee identification numbers or contact details in the questionnaires. Each completed questionnaire was assigned a random study code solely for data entry and statistical analysis.

The study was approved by the Ethics Committee of Le Van Thinh Hospital under Decision No. 09/HĐĐĐ-BVLVT, issued on June 28, 2024.

RESULTS

The median age of participants was 31 years, with an interquartile range of 8. The majority of healthcare workers were under 40 years old (85.4%). Females accounted for 70.3% of the participants. Nearly half of the study population was classified as overweight or obese. Regarding professional roles, most participants (62.5%) were non-physician healthcare workers (including nurses, midwives and technicians), while physicians accounted for 37.5% of the sample.

Regarding years of experience, most participants had worked in healthcare for 1 to less than 5 years (38.8%), followed by 5 to less than 10 years (26.7%) and 10 years or more (27.2%). The proportion of female participants was significantly higher among non-physician healthcare workers than among physicians ($p < 0.001$).

Table 1. Characteristics of the Study Population (N = 232)

Characteristic	Total, n (%) (N = 232)	Physicians, n (%) (N = 87)	Other HCWs, n (%) (N = 145)	p -value
Gender				
Male	69 (29.7)	44 (50.6)	25 (17.2)	<0.001
Female	163 (70.3)	43 (49.4)	120 (82.8)	
Body Mass Index (BMI)				
Underweight	14 (6.0)	10 (11.5)	4 (2.8)	0.064
Normal weight	115 (49.6)	38 (43.7)	77 (53.1)	
Overweight	46 (19.8)	20 (23.0)	26 (17.9)	
Obesity Class I	50 (21.6)	17 (19.5)	33 (22.8)	
Obesity Class II	7 (3.0)	2 (2.3)	5 (3.4)	
Age group (years)				
<30	92 (39.7)	41 (47.1)	51 (35.2)	0.315
30 – 39	106 (45.7)	35 (40.2)	71 (49.0)	
40 – 49	24 (10.3)	7 (8.0)	17 (11.7)	
50 – 59	10 (4.3)	4 (4.6)	6 (4.1)	
≥60	0 (0)	0 (0)	0 (0)	
Ethnicity				
Kinh	219 (94.4)	81 (93.1)	138 (95.2)	0.562
Other	13 (5.6)	6 (6.9)	7 (4.8)	
Years of medical experience				
<1	17 (7.3)	12 (13.8)	5 (3.4)	<0.001
1 – <5	90 (38.8)	43 (49.4)	47 (32.4)	
5 – <10	62 (26.7)	18 (20.7)	44 (30.3)	
≥10	63 (27.2)	14 (16.1)	49 (33.8)	

Table 2. Knowledge regarding obesity among healthcare workers (N = 232)

Survey Question	N (%)	95% CI
Perception of obesity		
1. Obesity is a disease	148 (63.8)	57.3-70.3%
2. Obesity is a health condition, but not a disease	84 (36.2)	29.7-42.7%
Knowledge of causes of obesity		
1. Nutritional causes	208 (89.7)	85.8-93.5%
2. Genetic causes	137 (59.1)	52.6-65.5%
3. Pathological causes	42 (18.1)	13.4-23.3%
4. Endocrine disorders	146 (62.9)	56.9-69.4%
5. Medication use	96 (41.4)	35.3-47.8%
6. Do not know	14 (6.0)	3.0-9.1%
Knowledge of obesity diagnostic methods		
1. Body Mass Index (BMI)	209 (90.1)	85.8-93.5%
2. Waist circumference measurement	127 (54.7)	48.3-61.2%
3. Abdominal ultrasound to assess subcutaneous fat	68 (29.3)	23.7-35.3%
4. Dual-energy X-ray absorptiometry (DEXA)	44 (19.0)	14.2-24.1%
5. Body fat–muscle index using specialized scales	111 (47.8)	40.9-54.3%
6. Do not know	21 (9.1)	5.6-12.5%
Knowledge of obesity treatment methods		
1. Dietary restriction	212 (91.4)	87.5-94.8%
2. Physical exercise	201 (86.6)	82.3-90.5%
3. Pharmacologic therapy	119 (51.3)	44.8-57.8%
4. Liposuction or fat-removal surgery	59 (25.4)	20.3-31.0%
5. Fat destruction using ultrasound	32 (13.8)	9.9-18.5%
6. Bariatric surgery (e.g., gastric reduction)	80 (34.5)	28.4-40.5%
7. Do not know	13 (5.6)	3.0-8.6%
Knowledge of anti-obesity medications approved by the Ministry of Health (Vietnam)		
1. Orlistat	115 (49.6)	43.1-56.0%
2. Sibutramine	23 (9.9)	6.5-13.8%
3. Liraglutide	76 (32.8)	27.2-38.8%
4. Diuretics	15 (6.5)	3.4-9.9%
5. Do not know	95 (40.9)	34.5-47.0%

Respondents could select multiple options; percentages exceed 100%.

Among non-physician healthcare workers, the proportion with less than one year of experience was the lowest. The remaining experience groups each accounted for approximately one-third of the sample. In contrast, among physicians, nearly 50% had 1 to less than 5 years of experience. This difference was statistically significant ($p < 0.001$).

There were no statistically significant differences between physicians and non-physician healthcare workers in BMI, age group or ethnicity. (Table 1)

Knowledge

Among the 232 participants, 148 (63.8%; 95% CI: 57.3-70.3%) considered obesity to be a disease. Nutritional factors were most commonly cited as the cause of obesity (89.7%; 95% CI: 85.8-93.5%), followed by endocrine disorders and genetic predisposition, each selected by over half of respondents.

Regarding methods for diagnosing obesity, more than 90% (90.1%; 95% CI: 85.8-93.5%) of participants identified body mass index (BMI) as the primary diagnostic tool, followed by waist circumference measurement (54.7%; 95% CI: 48.3-61.2%). Meanwhile, only 19.0% (95% CI: 14.2-24.1%) selected dual-energy X-ray absorptiometry (DEXA) and 47.8% (95% CI: 40.9-54.3%) chose body fat–muscle composition analysis using a specialized scale.

The two most commonly selected obesity treatment methods were dietary restriction (91.4%; 95% CI: 87.5-94.8%) and physical activity (86.6%; 95% CI: 82.3-90.5%). In contrast, only 51.3% (95% CI: 44.8-57.8%) selected pharmacologic treatment and 34.5% (95% CI: 28.4-40.5%) chose bariatric surgery.

Fewer than half of healthcare workers were aware of nationally approved pharmacotherapies (Orlistat 49.6% [95% CI: 43.1-56.0%], Liraglutide 32.8% [95% CI: 27.2-38.8%]), with 40.9% [95% CI: 34.5-47.0%] reporting no knowledge of possible treatments. (Table 2)

Attitudes

When asked about the roles of physicians and patients in obesity management, over 90% (90.1%; 95% CI: 86.2-93.5%) of participants agreed that physicians can help patients healthily lose weight, and 79.7% (95% CI: 75.0-84.9%) believed that the responsibility for weight loss lies with the patient. Regarding the initiation of obesity treatment, 93.5% (95% CI: 90.1-96.6%) of participants stated they would treat obesity in patients with comorbidities, while only 84.1% (95% CI: 79.3-88.4%) agreed to treat patients without comorbidities.

Regarding concern about obesity, more than half of the healthcare workers reported being very concerned about

Table 3. Attitudes and level of concern about obesity among healthcare workers

Survey Question	Total, n (%) (N = 232)	Physicians, n (%) (N = 87)	Other HCWs, n (%) (N = 145)	p-value
<i>“Physicians can help patients lose weight in a healthy way”</i>				
Agree	209 (90.1) [95% CI: 86.2-93.5%]	76 (87.4) [95% CI: 80.0-93.6%]	133 (91.7) [95% CI: 86.9-96.2%]	0.281*
Disagree	23 (9.9) [95% CI: 6.5-13.8%]	11 (12.6) [95% CI: 6.4-20.0%]	12 (8.3) [95% CI: 3.8-13.1%]	
<i>“Weight loss is the patient’s responsibility”</i>				
Agree	185 (79.7) [95% CI: 75.0-84.9%]	66 (75.9) [95% CI: 66.7-84.5%]	119 (82.1) [95% CI: 75.2-87.8%]	0.255*
Disagree	47 (20.3) [95% CI: 15.1-25.0%]	21 (24.1) [95% CI: 15.5-33.3%]	26 (17.9) [95% CI: 12.2-24.8%]	
<i>Would recommend obesity treatment if the patient has obesity with comorbidities</i>				
Agree	217 (93.5) [95% CI: 90.1-96.6%]	79 (90.8) [95% CI: 84.2-96.5%]	138 (95.2) [95% CI: 91.2-98.1%]	0.190*
Disagree	15 (6.5) [95% CI: 3.4-9.9%]	8 (9.2) [95% CI: 3.5-15.8%]	7 (4.8) [95% CI: 1.9-8.8%]	
<i>Would recommend obesity treatment if the patient has obesity without comorbidities</i>				
Agree	195 (84.1) [95% CI: 79.3-88.4%]	68 (78.2) [95% CI: 69.2-86.7%]	127 (87.6) [95% CI: 82.0-92.8%]	0.058*
Disagree	37 (15.9) [95% CI: 11.6-20.7%]	19 (21.8) [95% CI: 13.3-30.8%]	18 (12.4) [95% CI: 7.2-18.0%]	
<i>Level of concern about obesity in patients</i>				
Very concerned	126 (54.3) [95% CI: 47.8-60.3%]	47 (54.0) [95% CI: 43.2-64.4%]	79 (54.5) [95% CI: 45.9-62.7%]	0.971*
Somewhat concerned	86 (37.1) [95% CI: 31.0-43.1%]	32 (36.8) [95% CI: 26.7-46.7%]	54 (37.2) [95% CI: 29.5-45.4%]	
Not concerned	20 (8.6) [95% CI: 5.2-12.1%]	8 (9.2) [95% CI: 3.6-15.7%]	12 (8.3) [95% CI: 4.2-12.9%]	
*Pearson’s Chi-square test; **Fisher’s exact tests.				

*Pearson's Chi-square test; **Fisher's exact tests.

the issue, over one-third expressed some concern, and only 8.6% (95% CI: 5.2-12.1%) stated they were not concerned about obesity. There was no statistically significant difference in attitudes toward obesity or in the level of concern between physicians and other healthcare workers (Table 3).

Practices

In the management of patients with obesity, 83.6% (95% CI: 78.5-88.4%) of respondents recommended lifestyle modification for those with a BMI between 25 and 30 who had no comorbidities. For patients with a BMI between 30 and 35 without comorbidities, 72.4% (95% CI: 66.4-78.0%) recommended lifestyle modification, and 59.1% (95% CI: 52.6-65.5%) suggested referral to an obesity specialist. In patients with a BMI between 27 and 35 and comorbidities, three approaches were selected at nearly equal rates: lifestyle modification (66.4%; 95% CI: 59.9-72.0%), pharmacologic treatment (62.1%; 95% CI: 56.0-68.1%) and referral to an obesity specialty clinic (61.2%; 95% CI: 54.7-67.2%) (Table 4). Treatment recommendations varied depending on the patient's BMI and comorbidity profile, as illustrated in Figure 1.

When discussing barriers to communicating with patients about obesity, nearly 90% of healthcare workers reported encountering at least one obstacle. The most commonly cited barrier was “lack of patient motivation” (80.2%; 95% CI: 75.4-85.3%), followed by “patient sensitivity about weight” (57.3%; 95% CI: 50.9-63.4%). Other frequently reported barriers included “lack of time during consultations” (42.7%; 95% CI: 36.2-49.6%) and “lack of training in effective communication” (42.2%; 95% CI: 35.3-48.7%) (Table 4).

Compared with other healthcare workers, physicians were 2.3 times more likely to report time constraints during consultations ($p = 0.003$), 1.9 times more likely to cite insufficient healthcare resources ($p = 0.035$), and 4.5 times more likely to report inadequate knowledge of weight management

($p < 0.001$). The comparison of perceived barriers to discussing weight management between physicians and other health-care workers is summarized in Figure 2.

Regarding the influence of experience and BMI, stratified analyses revealed that healthcare workers with ≥ 5 years of experience demonstrated significantly higher rates of lifestyle modification recommendations for mild obesity compared to those with less experience ($p < 0.001$) (Table S1). However, there were no significant differences in KAP outcomes across participants' own BMI status (Table S2).

DISCUSSION

This study provides valuable insights into the knowledge, attitudes and practices of healthcare workers in Vietnam regarding obesity management. While general awareness of obesity as a health issue was relatively high, fewer than half of the participants were aware of approved pharmacotherapies. Notably, only 63.8% identified obesity as a chronic disease. This recognition rate is lower than those reported in recent Malaysian (88%), Swedish (91%) and broader ASEAN studies (84%), highlighting a persistent regional gap in understanding the metabolic complexity of obesity.¹⁴⁻¹⁷

Regarding attitudes, although 87.4% believed they could help patients lose weight, over three-quarters believed that weight loss is primarily the patient's responsibility. This perception is consistent with a prior Vietnamese study in which 55% of healthcare workers endorsed the same view.¹⁵ It is notably higher than the 41% reported in a regional study across South and Southeast Asia.¹⁴ The finding that 79.7% of healthcare workers attribute weight loss to patient responsibility reflects implicit weight bias widely documented in healthcare settings. This perspective is incongruent with current pathophysiological understanding of obesity as a chronic disease with strong genetic and neurobiological components. To improve

obesity care, training must address and normalize the biopsychosocial model of obesity.

Concern about obesity was not high among staff at Le Van Thinh Hospital; only 54.0% of physicians and 54.5% of other healthcare workers reported being highly concerned. To change entrenched and long-standing beliefs, a shift

toward a new public narrative grounded in modern scientific knowledge is needed.¹⁸

An interesting observation was the shift in treatment recommendations based on obesity severity. The declining frequency of lifestyle modification recommendations (from 83.6% for BMI 25–30 to 72.4% for BMI 30–35) despite

Table 4. Practices of healthcare workers regarding obesity management

Survey Question	Total, n (%) (N = 232)	Physicians, n (%) (N = 87)	Other HCWs, n (%) (N = 145)	p-value
Management of patients with BMI 25–30 and no comorbidities				
1. Lifestyle modification	194 (83.6) [95% CI: 78.5–88.4%]	74 (85.1) [95% CI: 77.1–92.6%]	120 (82.8) [95% CI: 75.9–88.7%]	0.647*
2. Pharmacologic treatment	25 (10.8) [95% CI: 6.9–15.5%]	9 (10.3) [95% CI: 3.8–17.0%]	16 (11.0) [95% CI: 6.3–17.0%]	0.870*
3. Referral to obesity specialist	47 (20.3) [95% CI: 15.1–26.3%]	19 (21.8) [95% CI: 13.3–30.6%]	28 (19.3) [95% CI: 12.9–26.3%]	0.643*
4. No intervention	14 (6.0) [95% CI: 3.0–9.1%]	0 (0.0) -	14 (9.7) [95% CI: 5.2–15.1%]	0.003*
5. Do not know what to do	25 (10.8) [95% CI: 6.9–15.1%]	10 (11.5) [95% CI: 4.8–18.4%]	15 (10.3) [95% CI: 5.8–15.6%]	0.785*
Management of patients with BMI 30–35 and no comorbidities				
1. Lifestyle modification	168 (72.4) [95% CI: 66.4–78.0%]	64 (73.6) [95% CI: 63.9–83.0%]	104 (71.7) [95% CI: 64.2–78.6%]	0.762*
2. Pharmacologic treatment	81 (34.9) [95% CI: 28.5–40.9%]	32 (36.8) [95% CI: 26.4–47.9%]	49 (33.8) [95% CI: 26.3–41.8%]	0.644*
3. Referral to obesity specialist	137 (59.1) [95% CI: 52.6–65.5%]	57 (65.5) [95% CI: 54.8–76.0%]	80 (55.2) [95% CI: 47.2–63.2%]	0.121*
4. No intervention	6 (2.6) [95% CI: 0.9–4.7%]	1 (1.1) [95% CI: 0.0–4.2%]	5 (3.4) [95% CI: 0.7–6.7%]	0.414**
5. Do not know what to do	31 (13.4) [95% CI: 9.1–17.7%]	8 (9.2) [95% CI: 3.4–15.7%]	23 (15.9) [95% CI: 10.3–21.9%]	0.149*
Management of patients with BMI 27–35 and comorbidities (e.g., HTN, T2DM, dyslipidemia)				
1. Lifestyle modification	154 (66.4) [95% CI: 59.9–72.0%]	54 (62.1) [95% CI: 51.7–73.2%]	100 (69.0) [95% CI: 61.5–76.5%]	0.282*
2. Pharmacologic treatment	144 (62.1) [95% CI: 56.0–68.1%]	54 (62.1) [95% CI: 51.7–72.0%]	90 (62.1) [95% CI: 53.9–69.7%]	1.000*
3. Referral to obesity specialist	142 (61.2) [95% CI: 54.7–67.2%]	65 (74.7) [95% CI: 65.3–83.7%]	77 (53.1) [95% CI: 45.1–61.4%]	0.001*
4. No intervention	3 (1.3) [95% CI: 0.0–3.0%]	0 (0.0) -	3 (2.1) [95% CI: 0.0–4.8%]	0.294**
5. Do not know what to do	30 (12.9) [95% CI: 8.6–17.2%]	8 (9.2) [95% CI: 3.4–15.7%]	22 (15.2) [95% CI: 9.7–21.2%]	0.189*
Barriers in discussing weight management with obese patients				
1. Lack of patient motivation	186 (80.2) [95% CI: 75.4–85.3%]	73 (83.9) [95% CI: 75.8–92.0%]	113 (77.9) [95% CI: 70.8–84.9%]	0.269*
2. Lack of time during consultation	99 (42.7) [95% CI: 36.2–49.6%]	48 (55.2) [95% CI: 43.2–65.9%]	51 (35.2) [95% CI: 27.7–42.3%]	0.003*
3. Patient sensitivity about weight	133 (57.3) [95% CI: 50.9–63.4%]	50 (57.5) [95% CI: 47.1–67.4%]	83 (57.2) [95% CI: 49.3–65.7%]	0.973*
4. Lack of primary care resources	69 (29.7) [95% CI: 23.7–35.3%]	33 (37.9) [95% CI: 27.3–48.3%]	36 (24.8) [95% CI: 17.9–32.2%]	0.035*
5. Lack of training in effective weight-related communication	98 (42.2) [95% CI: 35.3–48.7%]	43 (49.4) [95% CI: 38.3–59.8%]	55 (37.9) [95% CI: 30.1–46.3%]	0.086*
6. Physicians' lack of knowledge about weight management	37 (15.9) [95% CI: 11.2–20.7%]	25 (28.7) [95% CI: 19.3–38.0%]	12 (8.3) [95% CI: 4.1–12.8%]	<0.001*
7. No barriers reported	7 (3.0) [95% CI: 0.9–5.2%]	4 (4.6) [95% CI: 1.1–9.5%]	3 (2.1) [95% CI: 0.0–4.8%]	0.430**
8. Not applicable / never discussed obesity with patients	18 (7.8) [95% CI: 4.3–11.2%]	5 (5.7) [95% CI: 1.1–11.1%]	13 (9.0) [95% CI: 4.7–13.8%]	0.375*

Respondents could select multiple options; percentages exceed 100%. *Pearson's Chi-square test; **Fisher's exact tests.




Figure 1. Treatment recommendations (lifestyle, pharmacotherapy, referral) across BMI categories, with separate bars for physicians vs. non-physicians.




Figure 2. Forest plot of odds ratios and 95% CIs for barriers comparing physicians to other healthcare workers.

increasing metabolic dysfunction suggests potential therapeutic disbelief or uncertainty about the efficacy of pharmacotherapy. These findings align with studies from ASEAN countries, wherein, despite believing medications are effective, physicians rarely prescribe them.¹⁹ This may reflect a lack of pharmacotherapy knowledge, potentially correlating with low confidence in obesity treatment. Although multidisciplinary care models have proven effective, they remain underutilized in Vietnam.

When analyzing barriers, we found that physicians were significantly more likely than other hospital staff to report inadequate knowledge in weight management. Interestingly, physicians reported greater knowledge insufficiency (4.5x higher odds) than other staff. This may reflect greater awareness of knowledge gaps following more formal training, or increased time constraints limiting the opportunity to maintain currency with rapidly evolving obesity treatment guidelines. Barriers in counseling patients about obesity were considerable: 80% cited lack of patient motivation and 57.3% mentioned patient sensitivity about weight. Additionally, 28% of physicians reported insufficient knowledge as a major barrier, along with limited time and resources—factors more frequently reported by physicians than by other staff. Similar barriers have been reported across Asia. A review of 26 studies identified common challenges among healthcare providers, including a lack of knowledge, inadequate specialized training and insufficient systemic support.²⁰ Therefore, targeted interventions to enhance provider knowledge and skills are essential. Integrated training programs that combine medical knowledge, patient-centered communication and practical tools may help improve the quality of care and reduce delays in obesity management.

Our stratified analysis revealed a significant association between clinical experience and obesity management. Healthcare workers with five or more years of experience

demonstrated greater concern. They were more likely to recommend lifestyle modifications across various clinical scenarios than those with less than five years of experience. Notably, junior staff reported a significantly higher rate of uncertainty regarding appropriate interventions. This finding suggests that while fundamental knowledge may be acquired during formal education, the confidence and perceived importance of obesity management likely grow with clinical exposure and long-term patient interaction.

In contrast, the participants' own BMI status did not appear to significantly influence their knowledge, attitudes or clinical practices regarding obesity. This lack of association suggests that healthcare workers at Le Van Thinh Hospital generally maintain a professional approach to obesity care that is independent of their personal physical characteristics. This reinforces the idea that educational interventions and institutional protocols, rather than personal attributes, are the primary drivers of clinical decision-making in this setting.

To address these systemic and educational gaps, comprehensive system-level changes are required. Institutional interventions can include: (1) obesity medicine certification programs; (2) standardized screening and treatment protocols; (3) multidisciplinary care models with nutritionists, nurses, physiotherapists, and behavioral specialists; (4) electronic medical record (EMR) reminders for weight management; and (5) performance metrics linking obesity treatment to quality outcomes.

Although multidisciplinary care models have proven effective, they remain underutilized in Vietnam. To address this, an obesity specialty clinic was newly established at our institution in late 2023. However, within Vietnam's broader public hospital infrastructure, such dedicated units remain highly scarce. Obesity management is still mostly integrated into general endocrinology or nutrition

clinics rather than standalone centers. Furthermore, there is a lack of robust national data on the utilization rates at these specialized clinics, reflecting an urgent need for institutional investments in dedicated metabolic care models.

Limitations

This study has several limitations. First, the cross-sectional design reflects associations at a single point in time and does not establish causality among knowledge, attitudes and practices. Second, the use of convenience sampling during departmental meetings precludes the calculation of an exact response rate and may introduce selection bias. Although our sample predominantly consisted of nurses (53.9%) and internal medicine physicians (29.7%), which generally reflects the hospital's clinical workforce composition, some smaller clinical departments and surgical specialties may be under-represented, limiting the broader generalizability of the findings across all hospital sectors. Third, reliance on self-reported data may be subject to social desirability bias. Because our analysis involved multiple bivariate comparisons to explore associations between participant characteristics and various KAP domains, there is a potential for Type I error inflation. As no formal multiplicity adjustments were applied in this descriptive study, the statistically significant associations identified should be interpreted as exploratory and warrant confirmation in future adequately powered, hypothesis-driven research. While stratified analyses were performed to adjust for key confounders such as participant BMI and gender, full multivariable regression models were not utilized. The statistically significant associations identified in our bivariate comparisons should therefore be interpreted as exploratory. Finally, the study was conducted at a single hospital, which may not represent general trends across healthcare settings in Vietnam. Future research should include larger, geographically diverse samples and employ longitudinal designs to assess changes in perception and practice over time.

CONCLUSION

In conclusion, significant knowledge gaps and implicit weight biases persist among healthcare workers regarding obesity management. While this study was conducted at a single tertiary center, these findings highlight systemic educational barriers that likely extend across the national healthcare infrastructure. Addressing these gaps through specialized training, standardized protocols, and multi-disciplinary care models is not only critical for institutional improvement but also provides a vital framework for Vietnam and other ASEAN nations currently confronting similar rapid escalations in metabolic diseases.

Acknowledgments

The author sincerely thanks all healthcare workers who participated in the survey for their valuable time and insights. Special appreciation is extended to the Board of Directors and the heads of departments at Le Van Thinh Hospital for their support and facilitation in conducting this study.

Statement of Authorship

All authors certified fulfillment of ICMJE authorship criteria.

CRedit Author Statement

TT: Conceptualization, Methodology, Validation, Formal Analysis, Resources, Data curation, Writing – original draft preparation, Writing – review and editing.

Data Availability Statement

Datasets are not publicly available because participants in the study did not give written consent for their data to be shared.

Author Disclosure

The authors declared no conflict of interest.

Funding Source

None.

References

1. Jaacks LM, Vandevijvere S, Pan A, et al. The obesity transition: Stages of the global epidemic. *Lancet Diabetes Endocrinol.* 2019;7(3): 231-40. PMID: 30704950 PMCID: PMC7360432 DOI: 10.1016/S2213-8587(19)30026-9
2. World Health Organization. Obesity and overweight. Updated 2024. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
3. Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metabolism.* 2019;92:6-10. PMID: 30253139 DOI: 10.1016/j.metabol.2018.09.005
4. Bucher Della Torre S, Courvoisier DS, Saldarriaga A, Martin XE, Farpour-Lambert NJ. Knowledge, attitudes, representations and declared practices of nurses and physicians about obesity in a university hospital: training is essential. *Clin Obes.* 2018;8(2):122-30. PMID: 29327430 DOI: 10.1111/cob.12238
5. Siong TE, Florentino RF, Noor IM, Hlaing LM, Chotivichien S, Hop LT. A review of national plans of action for nutrition in Southeast Asian countries. *Malays J Nutr.* 2020;26(3):501-24. DOI: 10.31246/mjn-review-26-3
6. Nguyen TT, Maurizio T. Vietnam a country in transition: Health challenges. *BMJ Nutr Prev Health.* 2020;3(1):60-6. PMID: 33235972 PMCID: PMC7664505 DOI: 10.1136/bmjnp-2020-000069
7. Nguyen TT, Hoang MV. Non-communicable diseases, food and nutrition in Vietnam from 1975 to 2015: The burden and national response. *Asia Pac J Clin Nutr.* 2018;27(1):19-28. PMID: 29222878 DOI: 10.6133/apjcn.032017.13
8. Chu DT, Nguyet NTM, Dinh TC, et al. An update on physical health and economic consequences of overweight and obesity. *Diabetes Metab Syndr.* 2018;12(6):1095-100. PMID: 29799416 DOI: 10.1016/j.dsx.2018.05.004
9. Lauby-Secretan B, Scoccianti C, Loomis D, et al; International Agency for Research on Cancer Handbook Working Group. Body fatness and cancer -- Viewpoint of the IARC Working Group. *N Engl J Med.* 2016;375(8):794-8. PMID: 27557308 PMCID: PMC6754861 DOI: 10.1056/NEJMsrl606602
10. Phuc TT, Duc TQ, Quynh Chi VT, Quang PN. The prevalence of excess weight among Vietnamese adults: A pooled analysis of 58 studies with more than 430 thousand participants over the last three decades. *Nutr Health.* 2023;29(3):443-52. PMID: 36285514 DOI: 10.1177/02601060221129440
11. Tsukamoto Y, Wang H, Becker G, et al. Report of the Asian Forum of Chronic Kidney Disease Initiative (AFCKDI) 2007. "Current status and perspective of CKD in Asia": Diversity and specificity among Asian countries. *Clin Exp Nephrol.* 2009;13(3):249-56. PMID: 19288169 DOI: 10.1007/s10157-009-0156-8
12. Ministry of Health of Vietnam. Guidelines for diagnosis and management of obesity. Decision No. 2892/QĐ-BYT; 2022. <https://emohbackup.moh.gov.vn/publish/home?documentId=8975>
13. Dixon J, Hayden M, O'Brien P, Piterman L. Physician attitudes, beliefs and barriers towards the management and treatment of adult obesity: A literature review. *Aust J Prim Health.* 2008;14(3):9-18. DOI: 10.1071/PY08031
14. Tham KW, Ahmed A, Boonyavarakul A, et al. ACTION APAC: Understanding perceptions, attitudes and behaviours in obesity and its management across South and Southeast Asia. *Clin Obes.* 2024;14(3):e12644. PMID: 38332544 DOI: 10.1111/cob.12644
15. Tran NQ, Bich DNT, Nguyet TN, et al. Perceptions, attitudes, behaviours and barriers in obesity care: Findings from the ACTION-Vietnam study. *J ASEAN Fed Endocr Soc.* 2025;40(1):32-44. PMID: 40416487 PMCID: PMC12097987 DOI: 10.15605/jafes.040.01.17

16. Carrasco D, Thulesius H, Jakobsson U, Memarian E. Primary care physicians' knowledge and attitudes about obesity, adherence to treatment guidelines and association with confidence to treat obesity: A Swedish survey study. *BMC Prim Care*. 2022;23(1):208. PMID: 35971075 PMCID: PMC9378264 DOI: 10.1186/s12875-022-01811-x
17. Nor Hanipah Z, Abdul Ghani R, Goon MDME. ACTION Malaysia: Perception and barriers to obesity management among people with obesity and healthcare professionals in Malaysia. *BMC Public Health*. 2025;25(1):835. PMID: 40033266 PMCID: PMC11874754 DOI: 10.1186/s12889-025-22052-4
18. Rubino F, Puhl RM, Cummings DE, et al. Joint international consensus statement for ending stigma of obesity. *Nat Med*. 2020; 26(4):485-97. PMID: 32127716 PMCID: PMC7154011 DOI: 10.1038/s41591-020-0803-x
19. Boonyavarakul A, Buranapin S, Aniwat N, Chinthammit R. Perceptions, attitudes, and potential barriers to effective obesity care in Thailand: A survey of people with obesity and health care professionals from the ACTION APAC study. *J ASEAN Fed Endocr Soc*. 2023;38(Suppl 3):87. DOI: 10.15605/jafes.038.AFES.138
20. Anand VV, Zhe ELC, Chin YH, Lim WH, Goh RSJ, Lin C, et al. Barriers and facilitators to engagement with a weight management intervention in Asian patients with overweight or obesity: A systematic review. *Endocr Pract*. 2023;29(5):398-407. PMID: 36396016 DOI: 10.1016/j.eprac.2022.10.006

Authors are required to accomplish, sign and submit scanned copies of the JAFES Author Form consisting of: (1) Authorship Certification, that authors contributed substantially to the work, that the manuscript has been read and approved by all authors, and that the requirements for authorship have been met by each author; (2) the Author Declaration, that the article represents original material that is not being considered for publication or has not been published or accepted for publication elsewhere, that the article does not infringe or violate any copyrights or intellectual property rights; that no references have been made to predatory/suspected predatory journals; and that use of artificial intelligence (AI) or AI-assisted technologies shall be declared to include the name of the AI tool or service used; (3) the Author Contribution Disclosure, which lists the specific contributions of authors; (4) the Author Publishing Agreement which retains author copyright, grants publishing and distribution rights to JAFES, and allows JAFES to apply and enforce an Attribution-Non-Commercial Creative Commons user license; and (5) the Conversion to Visual Abstracts (*optional for original articles only) to improve dissemination to practitioners and lay readers. Authors are also required to accomplish, sign, and submit the signed ICMJE form for Disclosure of Potential Conflicts of Interest. For original articles, authors are required to submit a scanned copy of the Ethics Review Approval of their research as well as registration in trial registries as appropriate. For manuscripts reporting data from studies involving animals, authors are required to submit a scanned copy of the Institutional Animal Care and Use Committee approval. For Case Reports or Series, and Images in Endocrinology, consent forms, are required for the publication of information about patients; otherwise, appropriate ethical clearance has been obtained from the institutional review board. Articles and any other material published in the JAFES represent the work of the author(s) and should not be construed to reflect the opinions of the Editors or the Publisher.



Experience the new JAFES.
Visit us at www.ASEAN-endocrinejournal.org.

SUPPLEMENTARY TABLES

Table S1. Attitudes and practices stratified by years of experience (<5 years vs. ≥5 years)				
Survey question	Total, N (%)	<5 years experience, N (%)	≥5 years experience, N (%)	p-value
<i>“Physicians can help patients lose weight in a healthy way”</i>				
Agree	209 (90.1) [95% CI: 86.2-93.5%]	93 (86.9) [95% CI: 80.5-93.3%]	116 (92.8) [95% CI: 88.0-96.9%]	0.135*
Disagree	23 (9.9) [95% CI: 6.5-13.8%]	14 (13.1) [95% CI: 6.7-19.5%]	9 (7.2) [95% CI: 3.1-12.0%]	
<i>“Weight loss is the patient’s responsibility”</i>				
Agree	185 (79.7) [95% CI: 75.0-84.9%]	82 (76.6) [95% CI: 68.6-84.4%]	103 (82.4) [95% CI: 74.8-89.4%]	0.276*
Disagree	47 (20.3) [95% CI: 15.1-25.0%]	25 (23.4) [95% CI: 15.6-31.4%]	22 (17.6) [95% CI: 10.6-25.2%]	
<i>Would recommend obesity treatment if the patient has obesity with comorbidities</i>				
Agree	217 (93.5) [95% CI: 90.1-96.6%]	97 (90.7) [95% CI: 85.0-96.3%]	120 (96.0) [95% CI: 92.2-99.2%]	0.099*
Disagree	15 (6.5) [95% CI: 3.4-9.9%]	10 (9.3) [95% CI: 3.7-15.0%]	5 (4.0) [95% CI: 0.8-7.8%]	
<i>Would recommend obesity treatment if the patient has obesity without comorbidities</i>				
Agree	195 (84.1) [95% CI: 79.3-88.4%]	84 (78.5) [95% CI: 70.8-86.5%]	111 (88.8) [95% CI: 83.3-94.6%]	0.033*
Disagree	37 (15.9) [95% CI: 11.6-20.7%]	23 (21.5) [95% CI: 13.5-29.2%]	14 (11.2) [95% CI: 5.4-16.7%]	
<i>Level of concern about obesity in patients</i>				
Very concerned	126 (54.3) [95% CI: 47.8-60.3%]	50 (46.7) [95% CI: 37.3-56.4%]	76 (60.8) [95% CI: 51.1-69.3%]	0.027*
Somewhat concerned	86 (37.1) [95% CI: 31.0-43.1%]	43 (40.2) [95% CI: 31.1-49.5%]	43 (34.4) [95% CI: 26.4-42.7%]	
Not concerned	20 (8.6) [95% CI: 5.2-12.1%]	14 (13.1) [95% CI: 6.4-19.6%]	6 (4.8) [95% CI: 1.5-8.8%]	
<i>Management of patients with BMI 25–30 and no comorbidities</i>				
1. Lifestyle modification	194 (83.6) [95% CI: 78.5-88.4%]	79 (73.8) [95% CI: 65.4-82.2%]	115 (92.0) [95% CI: 86.8-96.3%]	<0.001*
2. Pharmacologic treatment	25 (10.8) [95% CI: 6.9-15.5%]	11 (10.3) [95% CI: 5.0-16.8%]	14 (11.2) [95% CI: 6.0-17.1%]	0.822*
3. Referral to obesity specialist	47 (20.3) [95% CI: 15.1-26.3%]	13 (12.1) [95% CI: 5.6-18.9%]	34 (27.2) [95% CI: 19.5-35.0%]	0.004*
4. No intervention	14 (6.0) [95% CI: 3.0-9.1%]	7 (6.5) [95% CI: 2.4-11.6%]	7 (5.6) [95% CI: 1.8-12.0%]	0.764*
5. Do not know what to do	25 (10.8) [95% CI: 6.9-15.1%]	20 (18.7) [95% CI: 11.3-26.7%]	5 (4.0) [95% CI: 0.8-7.8%]	<0.001*
<i>Management of patients with BMI 30–35 and no comorbidities</i>				
1. Lifestyle modification	168 (72.4) [95% CI: 66.4-78.0%]	71 (66.4) [95% CI: 57.8-75.4%]	97 (77.6) [95% CI: 69.6-84.7%]	0.056*
2. Pharmacologic treatment	81 (34.9) [95% CI: 28.5-40.9%]	40 (37.4) [95% CI: 28.4-46.9%]	41 (32.8) [95% CI: 24.1-41.0%]	0.465*
3. Referral to obesity specialist	137 (59.1) [95% CI: 52.6-65.5%]	60 (56.1) [95% CI: 46.5-65.7%]	77 (61.6) [95% CI: 52.7-70.2%]	0.394*
4. No intervention	6 (2.6) [95% CI: 0.9-4.7%]	2 (1.9) [95% CI: 0.0-4.8%]	4 (3.2) [95% CI: 0.7-6.6%]	0.689**
5. Do not know what to do	31 (13.4) [95% CI: 9.1-17.7%]	19 (17.8) [95% CI: 10.4-25.0%]	12 (9.6) [95% CI: 4.7-15.3%]	0.069*
<i>Management of patients with BMI 27–35 and comorbidities (e.g., HTN, T2DM, dyslipidemia)</i>				
1. Lifestyle modification	154 (66.4) [95% CI: 60.3-72.8%]	60 (56.1) [95% CI: 46.5-65.1%]	94 (75.2) [95% CI: 67.4-83.2%]	0.002*
2. Pharmacologic treatment	144 (62.1) [95% CI: 56.0-68.1%]	61 (57.0) [95% CI: 47.7-66.1%]	83 (66.4) [95% CI: 57.6-75.0%]	0.142*
3. Referral to obesity specialist	142 (61.2) [95% CI: 54.7-67.2%]	66 (61.7) [95% CI: 52.8-71.4%]	76 (60.8) [95% CI: 51.9-69.2%]	0.891*
4. No intervention	3 (1.3) [95% CI: 0.0-3.0%]	1 (0.9) [95% CI: 0.0-3.0%]	2 (1.6) [95% CI: 0.0-3.9%]	1.000**
5. Do not know what to do	30 (12.9) [95% CI: 8.6-17.2%]	19 (17.8) [95% CI: 10.4-25.0%]	11 (8.8) [95% CI: 4.0-14.0%]	0.043*
<i>Barriers in discussing weight management with obese patients</i>				
1. Lack of patient motivation	186 (80.2) [95% CI: 75.4-85.3%]	85 (79.4) [95% CI: 71.3-87.1%]	101 (80.8) [95% CI: 73.8-87.6%]	0.796*
2. Lack of time during consultation	99 (42.7) [95% CI: 36.2-49.6%]	50 (46.7) [95% CI: 37.7-55.6%]	49 (39.2) [95% CI: 30.5-48.1%]	0.248*
3. Patient sensitivity about weight	133 (57.3) [95% CI: 50.9-63.4%]	60 (56.1) [95% CI: 46.9-65.4%]	73 (58.4) [95% CI: 49.5-66.7%]	0.721*
4. Lack of primary care resources	69 (29.7) [95% CI: 23.7-35.3%]	32 (29.9) [95% CI: 21.4-38.7%]	37 (29.6) [95% CI: 21.9-37.8%]	0.959*
5. Lack of training in effective weight-related communication	98 (42.2) [95% CI: 35.3-48.7%]	46 (43.0) [95% CI: 34.2-52.6%]	52 (41.6) [95% CI: 32.5-50.4%]	0.831*
6. Physicians' lack of knowledge about weight management	37 (15.9) [95% CI: 11.2-20.7%]	18 (16.8) [95% CI: 10.1-24.3%]	19 (15.2) [95% CI: 8.8-21.3%]	0.737*
7. No barriers reported	7 (3.0) [95% CI: 0.9-5.2%]	3 (2.8) [95% CI: 0.0-6.3%]	4 (3.2) [95% CI: 0.8-6.6%]	1.000**
8. Not applicable / never discussed obesity with patients	18 (7.8) [95% CI: 4.3-11.2%]	10 (9.3) [95% CI: 4.0-15.3%]	8 (6.4) [95% CI: 2.3-11.4%]	0.403*
Respondents could select multiple options: percentages exceed 100%. *Pearson’s Chi-square test; **Fisher’s exact tests.				

Table S2. Relationship between participant BMI status and obesity attitudes/practices

Survey question	Total, N (%)	Underweight / normal weight HCWs, N (%)	Overweight / obese HCWs, N (%)	p-value
<i>“Physicians can help patients lose weight in a healthy way”</i>				
Agree	209 (90.1) [95% CI: 86.2-93.5%]	115 (89.1) [95% CI: 83.7-94.7%]	94 (91.3) [95% CI: 84.8-96.4%]	0.592*
Disagree	23 (9.9) [95% CI: 6.5-13.8%]	14 (10.9) [95% CI: 5.3-16.3%]	9 (8.7) [95% CI: 3.6-15.2%]	
<i>“Weight loss is the patient’s responsibility”</i>				
Agree	185 (79.7) [95% CI: 75.0-84.9%]	102 (79.1) [95% CI: 72.1-85.7%]	83 (80.6) [95% CI: 72.3-88.0%]	0.776*
Disagree	47 (20.3) [95% CI: 15.1-25.0%]	27 (20.9) [95% CI: 14.3-27.9%]	20 (19.4) [95% CI: 12.0-27.7%]	
<i>Would recommend obesity treatment if the patient has obesity with comorbidities</i>				
Agree	217 (93.5) [95% CI: 90.1-96.6%]	121 (93.8) [95% CI: 89.4-97.7%]	96 (93.2) [95% CI: 88.0-97.7%]	0.855*
Disagree	15 (6.5) [95% CI: 3.4-9.9%]	8 (6.2) [95% CI: 2.3-10.6%]	7 (6.8) [95% CI: 2.3-12.0%]	
<i>Would recommend obesity treatment if the patient has obesity without comorbidities</i>				
Agree	195 (84.1) [95% CI: 79.3-88.4%]	105 (81.4) [95% CI: 74.4-88.1%]	90 (87.4) [95% CI: 80.7-94.0%]	0.216*
Disagree	37 (15.9) [95% CI: 11.6-20.7%]	24 (18.6) [95% CI: 11.9-25.6%]	13 (12.6) [95% CI: 6.0-19.3%]	
<i>Level of concern about obesity in patients</i>				
Very concerned	126 (54.3) [95% CI: 47.8-60.3%]	65 (50.4) [95% CI: 41.6-58.8%]	61 (59.2) [95% CI: 50.0-68.6%]	0.365*
Somewhat concerned	86 (37.1) [95% CI: 31.0-43.1%]	51 (39.5) [95% CI: 31.6-48.8%]	35 (34.0) [95% CI: 24.8-43.2%]	
Not concerned	20 (8.6) [95% CI: 5.2-12.1%]	13 (10.1) [95% CI: 5.1-16.0%]	7 (6.8) [95% CI: 2.1-12.4%]	
<i>Management of patients with BMI 25–30 and no comorbidities</i>				
1. Lifestyle modification	194 (83.6) [95% CI: 78.5-88.4%]	104 (80.6) [95% CI: 74.0-87.0%]	90 (87.4) [95% CI: 80.9-93.8%]	0.167*
2. Pharmacologic treatment	25 (10.8) [95% CI: 6.9-15.5%]	12 (9.3) [95% CI: 4.5-14.2%]	13 (12.6) [95% CI: 6.9-19.1%]	0.418*
3. Referral to obesity specialist	47 (20.3) [95% CI: 15.1-26.3%]	25 (19.4) [95% CI: 12.6-26.8%]	22 (21.4) [95% CI: 13.7-29.0%]	0.709*
4. No intervention	14 (6.0) [95% CI: 3.0-9.1%]	9 (7.0) [95% CI: 3.1-11.1%]	5 (4.9) [95% CI: 1.1-9.4%]	0.500*
5. Do not know what to do	25 (10.8) [95% CI: 6.9-15.1%]	18 (14.0) [95% CI: 8.2-20.7%]	7 (6.8) [95% CI: 2.0-11.9%]	0.081*
<i>Management of patients with BMI 30–35 and no comorbidities</i>				
1. Lifestyle modification	168 (72.4) [95% CI: 66.4-78.0%]	90 (69.8) [95% CI: 60.8-77.5%]	78 (75.7) [95% CI: 67.3-84.0%]	0.313*
2. Pharmacologic treatment	81 (34.9) [95% CI: 28.5-40.9%]	47 (36.4) [95% CI: 28.3-44.4%]	34 (33.0) [95% CI: 24.7-42.3%]	0.587*
3. Referral to obesity specialist	137 (59.1) [95% CI: 52.6-65.5%]	78 (60.5) [95% CI: 51.6-68.6%]	59 (57.3) [95% CI: 47.6-66.7%]	0.624*
4. No intervention	6 (2.6) [95% CI: 0.9-4.7%]	3 (2.3) [95% CI: 0.0-5.3%]	3 (2.9) [95% CI: 0.0-6.6%]	1.000**
5. Do not know what to do	31 (13.4) [95% CI: 9.1-17.7%]	21 (16.3) [95% CI: 10.4-23.3%]	10 (9.7) [95% CI: 4.0-15.5%]	0.144*
<i>Management of patients with BMI 27–35 and comorbidities (e.g., HTN, T2DM, dyslipidemia)</i>				
1. Lifestyle modification	154 (66.4) [95% CI: 60.3-72.8%]	86 (66.7) [95% CI: 57.6-74.8%]	68 (66.0) [95% CI: 56.1-75.5%]	0.917*
2. Pharmacologic treatment	144 (62.1) [95% CI: 56.0-68.1%]	81 (62.8) [95% CI: 53.3-70.8%]	63 (61.2) [95% CI: 51.5-70.2%]	0.800*
3. Referral to obesity specialist	142 (61.2) [95% CI: 54.7-67.2%]	77 (59.7) [95% CI: 51.2-67.8%]	65 (63.1) [95% CI: 53.1-72.5%]	0.596*
4. No intervention	3 (1.3) [95% CI: 0.0-3.0%]	2 (1.6) [95% CI: 0.0-4.1%]	1 (1.0) [95% CI: 0.0-3.3%]	1.000**
5. Do not know what to do	30 (12.9) [95% CI: 8.6-17.2%]	21 (16.3) [95% CI: 10.2-22.9%]	9 (8.7) [95% CI: 3.5-14.1%]	0.089*
<i>Barriers in discussing weight management with obese patients</i>				
	232 (100.0)	129 (100.0)	103 (100.0)	
1. Lack of patient motivation	186 (80.2) [95% CI: 75.4-85.3%]	104 (80.6) [95% CI: 73.2-86.9%]	82 (79.6) [95% CI: 71.3-87.2%]	0.848*
2. Lack of time during consultation	99 (42.7) [95% CI: 36.2-49.6%]	54 (41.9) [95% CI: 33.9-50.8%]	45 (43.7) [95% CI: 34.0-53.3%]	0.780*
3. Patient sensitivity about weight	133 (57.3) [95% CI: 50.9-63.4%]	77 (59.7) [95% CI: 51.5-68.5%]	56 (54.4) [95% CI: 44.9-63.2%]	0.416*
4. Lack of primary care resources	69 (29.7) [95% CI: 23.7-35.3%]	43 (33.3) [95% CI: 24.6-41.3%]	26 (25.2) [95% CI: 16.8-34.4%]	0.180*
5. Lack of training in effective weight-related communication	98 (42.2) [95% CI: 35.3-48.7%]	56 (43.4) [95% CI: 34.5-52.4%]	42 (40.8) [95% CI: 31.7-50.5%]	0.687*
6. Physicians' lack of knowledge about weight management	37 (15.9) [95% CI: 11.2-20.7%]	26 (20.2) [95% CI: 13.2-27.5%]	11 (10.7) [95% CI: 5.1-17.4%]	0.050*
7. No barriers reported	7 (3.0) [95% CI: 0.9-5.2%]	4 (3.1) [95% CI: 0.7-6.6%]	3 (2.9) [95% CI: 0.0-6.4%]	1.000**
8. Not applicable / never discussed obesity with patients	18 (7.8) [95% CI: 4.3-11.2%]	11 (8.5) [95% CI: 4.2-13.7%]	7 (6.8) [95% CI: 2.2-11.9%]	0.624*
Respondents could select multiple options; percentages exceed 100%. *Pearson's Chi-square test; **Fisher's exact tests.				